

A MODULI FIXING MECHANISM IN M THEORY.B.S. ACHARYA¹

Department of Physics,
Rutgers University,
126 Freylinghuysen Road,
NJ 08854-0849.

ABSTRACT

We study M theory compactifications on manifolds of G_2 -holonomy with gauge and matter fields supported at singularities. We show that, under certain topological conditions, the combination of background G -flux and background fields at the singularities induces a potential for the moduli with an isolated minimum. The theory in the minimum is supersymmetric and has a negative cosmological constant in the simplest case. In a more realistic scenario, we find that the fundamental scale is around 10 Tev and the hierarchy between the four dimensional Planck and electroweak scales may be explained by the value of a topological invariant. Hyperbolic three-manifolds enter the discussion in an interesting way.

¹bacharya@physics.rutgers.edu

1. Introduction.

One of the traditional problems which has plagued string theory and subsequently M theory is the lack of predictivity of the low energy effective action. There are apparently many vacua of the theory. For example, in vacua with compact extra dimensions, there are many ways to choose the compactification data. Each such choice describes different physics and most of these choices do not seem to describe anything resembling physics as we know it. At present we know of no compelling reason why the theory might favour universes with three spatial dimensions let alone a local symmetry group $SU(3) \times SU(2) \times U(1)$ with the elementary particles living in three complex representations. On the other hand, we presently don't know of any other non-anthropocentric explanations of these facts either, so string theory is no better than any other theory in this regard.

A simple manifestation of the vacuum degeneracy/predictivity problem is that string and M theory compactifications have moduli. At the classical level this is due to the fact that the low energy effective potential has many flat directions and the values of scalar fields which determine the size and shape of the extra dimensions remain undetermined by the theory. At the quantum level there is a similar sort of problem in that non-perturbative effects such as instantons appear to generate a potential which pushes the fields off to extreme weak coupling and large volume [1]. In this regime, the potential is approximately flat again.

Naively one might think that the runaway potential might be useful for inflation or quintessence but the non-perturbative potentials that arise this way apparently do not inflate [2].

In this paper we will study M theory compactifications to four dimensions on manifolds of G_2 -holonomy. Many such compactifications arise as a certain limit of Calabi-Yau compactifications of the heterotic string. Calabi-Yau compactifications have traditionally been regarded as the most promising for obtaining realistic particle physics from string theory. The limit we will study is not very well approximated by weakly coupled heterotic strings and is much better described by M theory on a compact manifold X of G_2 -holonomy. Note that X must have certain kinds of singularities in order for the low energy dynamics to be described by a non-Abelian gauge theory coupled to chiral fermions. The singularities leading to non-Abelian gauge groups were first studied in [3] and in more detail in [4, 5, 6] whereas singularities which support chiral fermions have been studied in [6, 7, 8, 9]. Applications of these studies to more detailed questions of phenomenology have been made in [10, 11].

In particular, in order to obtain non-Abelian local symmetries, the G_2 -

manifold X contains a three manifold Q of orbifold singularities [3]. Near Q , X looks like $\mathbf{R}^4/\Gamma \times Q$. Hence, X has an orbifold singularity along Q . Γ is a finite subgroup of $SU(2)$ acting as a subgroup of $SO(4)$ on $\mathbf{R}^4$. Quarks and leptons emerge from additional singularities of X through which Q passes.

Up until relatively recently, most studies of string compactification focussed on vacua without fluxes. Typically the metric (as well as non-Abelian gauge fields in the case of heterotic strings) is the only non-zero background field and one studies the theory around a minimum of the classical potential. Our comments above on flat directions in the classical potential apply to these cases. More recently however, attention has been devoted to the study of compactifications with flux. These are compactifications in which the various p -form gauge fields present in the theory have been activated: the theory is studied in the presence of non-zero electric and/or magnetic fields. Since the kinetic energy of the p -form field depends upon the space-time metric, which in turn depends upon the moduli, flux compactifications lead to a potential for the moduli. Moduli stabilisation due to fluxes has been studied in a variety of papers [12] – [30]. An open question is can this approach lead to a potential which fixes all the moduli in a compactification to four dimensions?

In the context of G_2 -compactifications, the flux induced potential is positive [30] and so also suffers from a runaway problem. Since most of the interesting physics of G_2 -compactifications involves the fields localised at singularities a simple question that one can ask is can this runaway be stabilised by turning on the fields supported at singularities of the G_2 -manifold? We will study this question in this paper.

We will be interested in the sorts of singular G_2 -manifolds that give rise to Yang-Mills theory and chiral fermions and will study the effect of turning on these fields which are supported at the singularities. We will show that, as long as a certain topological invariant, c_2 , is non-zero and suitably large, that the combination of both fluxes and the fields supported at singularities is enough to fix all the moduli of the G_2 -manifold.

The size of the G_2 -manifold in the minimum of the potential is large if c_2 is large, so that the approximation to the potential that we use is valid. Thus, if this is indeed the case then these minima produce four dimensional vacua of M theory with no undetermined parameters. All masses and couplings are in principle calculable by the theory, although this is difficult in practice.

In the next section we describe the equations describing the absolute minimum of the supergravity potential and explain how they can be solved and why the solution is isolated. An important role is played by a certain complex Chern-Simons invariant $c_1 + ic_2$ which gives an additive contribution to the superpotential. We show that c_2 is non-zero and can be large when Q

is a hyperbolic three manifold.

The moduli fixing mechanism presented here involves only the gauge fields along 3-manifolds Q (and their superpartners) and as such can be applied to *non-realistic* G_2 -compactifications which do not include quarks and leptons. In this case, the vacuum we describe is supersymmetric and has a negative cosmological constant Λ_0 . In the concluding section, we consider the possibility that the moduli fixing mechanism is useful in a more realistic context. For this it is necessary to assume that the singularities of X support also a supersymmetric standard model *and* a supersymmetry breaking sector. Since by assumption supersymmetry is broken, particle physics interactions will generate a contribution Λ_1 to the cosmological constant and the main idea is to balance Λ_0 against Λ_1 . This is necessary if one is to obtain (at the very least) a positive cosmological constant.

In this scenario, we find that the fundamental Planck scale is 10 Tev (or smaller) and that the much higher four dimensional Planck scale emerges because the topological invariant c_2 is large. Thus, the heirarchy of scales that we observe between particle physics and gravity in four dimensions might be explained in these models by the fact that a topological invariant takes the value that it does.

2. Vacuum Equations for Moduli.

M theory on a compact G_2 -manifold X has a low energy description in terms of a minimally supersymmetric supergravity theory in four dimensions. This theory is coupled to $b_3(X)$ moduli superfields. If X contains a 3-manifold Q along which there is an orbifold singularity then there will also be non-Abelian gauge fields. Note that there could be several different three manifolds Q_a of orbifold singularities so there can be more than one gauge group factor. If Q_a passes through certain additional conical singularities there can be chiral superfields in a complex representation of the gauge group associated to Q_a .

For the purposes of explaining the simple mechanism which fixes the moduli we will only require that there is a single gauge group and therefore a single Q . We wont need to talk about the matter fields. The classical solution to the low energy equations of motion is a solution of eleven dimensional supergravity in which the only non-zero field is the 11-metric. This is a product of the G_2 -holonomy metric on X and flat four dimensional Minkowski spacetime. Since the superpotential is exactly zero in this classical low energy limit, the potential for the moduli fields is also exactly zero.

We will ask the simple question: what happens to the theory if we turn on a background four-form field G and a background gauge field A along Q ?

In the case with $A = 0$, this question has already been studied. The answer is that G induces a superpotential which is linear in the moduli superfields. In this case, the resulting potential for the moduli is positive and runs off to zero at infinity [30]. This corresponds to the large volume limit.

As we will explain momentarily, turning on the gauge field and its bosonic superpartners leads to a modification of the superpotential, which in the simplest non-trivial possibility is via the addition of a constant, $c_1 + ic_2$. This constant, or rather its imaginary part, has a dramatic effect on the potential: its presence implies a supersymmetric minimum with negative cosmological constant. Moreover, all the moduli fields are fixed in the minimum. In the remainder of this section we will explain these results in more detail. In order to do that we need to provide some basic background material. A more detailed explanation of some of the mathematics may be found in [31].

2.1. Kaluza-Klein on G_2 -manifolds.

A manifold of G_2 -holonomy always comes equipped with a 3-form φ which is covariantly constant (parallel) in the G_2 -holonomy metric. The existence of φ is equivalent to the existence of a G_2 -holonomy metric. Given a fixed G_2 -holonomy metric on a compact 7-manifold X , a new metric with the same holonomy can be constructed by perturbing φ by any small harmonic three-form. Thus the moduli space of compact G_2 -manifolds is locally isomorphic to a small neighbourhood of a point in $H^3(X, \mathbf{R})$. There are thus $b_3(X)$ moduli fields in the four dimensional effective action. These combine with harmonic fluctuations of the 3-form potential C of low energy M theory to give $b_3(X)$ complex scalars which become the lowest components of scalar superfields in four dimensions.

In more detail, let $\phi_j(x)$ $j = 1, \dots, b_3$ be a basis for $H^3(X, \mathbf{R})$. Then the Kaluza-Klein harmonic ansatz is

$$C + i\varphi = \sum_j Z_j(y) \phi_j(x) \quad (1)$$

The $Z_j(y)$'s are complex scalar fields in four dimensions and y denotes coordinates of four dimensional spacetime. We define the real and imaginary parts of the complex scalars by

$$Z_j = t_j + is_j \quad (2)$$

The t_j 's are axions because of C -field gauge transformations. The s_j 's are the moduli fields whose vacuum expectation values determine the size and shape of the G_2 -holonomy metric on X . This is because changing the s_j 's changes φ and therefore also the metric on X .

As argued in [15, 29] and proven in [30], turning on a background field strength G for C induces the following superpotential into the four dimensional theory

$$W = \frac{1}{8\pi^2} \int_X \left(\frac{C}{2} + i\varphi \right) \wedge G \quad (3)$$

Note that as explained in [30] the real part of W is classically an angular variable of period $\frac{1}{2}$. In the quantum theory, to account for the effects of G -flux through the four-dimensional spacetime, we should accept all possible values of $\text{Re}W$ which differ by $\frac{N}{2}$, for all integers N . Although important, the vacuum states distinguished by N will not figure much in what follows.

Since G does not depend upon the metric and is harmonic and quantised, we may expand it in terms of a basis of harmonic 4-forms ρ_j which are dual to the ϕ_j . This means that the superpotential can be written as

$$W = \frac{1}{8\pi^2} \Sigma_k Z_k G_k \quad (4)$$

and is therefore linear in the moduli superfields with coefficients which are the flux quanta².

Finally, since we will be interested in general supersymmetric vacua, we will require the Kahler potential and its derivatives. The Kahler potential depends only on the s_k 's and not on the axions. It is given by [32, 33, 30]

$$K = -3 \ln \left(a \int_X \varphi \wedge * \varphi \right) \quad (5)$$

where a is $\frac{1}{14\pi^2}$ and $*\varphi$ is the 4-form which is Hodge dual of φ in the metric on X . Note that since $*$ depends nonlinearly on the metric, e^K is a nonlinear function of the moduli. The first derivative of the Kahler potential with respect to the Z_k is [31, 30]

$$\frac{\partial K}{\partial Z_k} = \frac{7}{2} i \frac{\int_X \phi_k \wedge * \varphi}{\int_X \varphi \wedge * \varphi} \quad (6)$$

2.2. Superpotential for Gauge Fields.

In this subsection we will derive the contribution to the superpotential from the gauge fields and their superpartners which are localised along Q (times four dimensional spacetime $M^{3,1}$). When the eleven dimensional spacetime has an A , D or E orbifold singularity along a 7-manifold Y such

²The quanta G_k are integer multiples of 2π .

as $Q \times M^{3,1}$, as long as Y is large compared to the Planck length, there is a description of the M theory physics near Y in terms of seven dimensional super Yang-Mills theory on Y with an A, D, E gauge group G determined by the orbifold singularity [3]. The bosonic fields of 7d super Yang-Mills in flat space are a gauge field A and three scalars B . In our context, the Yang-Mills theory is on a curved spacetime, but because Q is a supersymmetric three cycle in a manifold of G_2 -holonomy the classical description of the physics for large Q is still supersymmetric even though the spacetime is curved³. Of the sixteen supersymmetries present in the flat space gauge theory, only four remain once we compactify on Q . For large Q there is then a description of the dynamics in terms of a supersymmetric Yang-Mills theory on M with minimal supersymmetry.

An important point is that the three scalars B actually become the components of a 1-form on Q because of supersymmetry. A very simple way to see this is that if we consider the fluctuations of the three components of A which are along Q , then these fluctuations look like scalar fields in four dimensions. But a real scalar field in four dimensional supersymmetry must combine with either a 2-form potential or another scalar field in order to form a representation of the supersymmetry algebra. In this case the three components of B transform as 1-forms on Q and not as 2-forms on M [3]. Thus the components of A tangential to Q have bosonic superpartners which are a 1-form B on Q . All fields transform in the adjoint representation of the gauge group.

One can check explicitly (see appendix) that the conditions for unbroken supersymmetry imply that

$$F(A) = B \wedge B \quad (7)$$

and

$$D_A B = 0 \quad (8)$$

ie that the field strength of the gauge field components along Q are equal to the commutator of the B 's and that the covariant derivative of B with respect to A is zero.

These equations are the critical points of the following complex functional

$$\omega \equiv \int_Q \text{tr}(A + iB) \wedge d(A + iB) + \frac{2}{3}(A + iB) \wedge (A + iB) \wedge (A + iB) \quad (9)$$

with respect to the complex gauge field $A + iB$. ω is the complex Chern-Simons functional. Its critical points are complex flat connections. These

³See [3] for a detailed discussion of supersymmetry and a description of the low energy physics in this context.

are equivalent to a representation of the fundamental group of Q in the complexification of the ADE gauge group $G^{\mathbb{C}}$ modulo conjugation.

This implies that the Chern-Simons functional ω is the contribution to the superpotential of the four dimensional theory from fluctuations of A and B along Q . ω is invariant under gauge transformations of the complexification of the gauge group as F -terms should be. In fact we can use the latter fact to rederive ω :

The contribution to the four dimensional potential energy coming from the fluctuations of A and B along Q is given by the bosonic part of the super Yang-Mills action on Q . This includes the Yang-Mills action for A

$$V = \int_Q \text{tr} F(A) \wedge *F(A) + \dots \quad (10)$$

Then we can ask, what is the superpotential W for which

$$V = \left| \frac{\delta W}{\delta(A + iB)} \right|^2 \quad (11)$$

Since the real Chern-Simons functional, $\omega_{\mathbf{R}}$ for A (which is ω with $B = 0$) has the property that

$$\delta\omega_{\mathbf{R}} \propto F(A) \quad (12)$$

complex gauge invariance of the F -term uniquely fixes ω to be the complex Chern-Simons functional⁴

As an aside, we note that for G_2 -compactifications which are dual to heterotic or Type I or other IIB orientifold compactifications on Calabi-Yau threefolds, ω is the M theory ‘dual’ of the holomorphic Chern-Simons functional which is the superpotential in these string compactifications.

We are going to pick our background gauge field $A_o + iB_o$ to simply be a critical point of ω ie a flat complex gauge field. We are also going to assume that the moduli space of complex flat connections is zero dimensional ie that the flat connection we choose is isolated. This means from the low energy four dimensional point of view that all the fluctuations of the gauge field around this background are massive. If this were not the case then the low energy theory would contain phenomenologically undesirable massless adjoint fields and the Kahler potential above would also require modification to include these additional light degrees of freedom.

The Chern-Simons functional ω is a topologically invariant functional. In particular, its value on a background flat connection is a constant

$$\omega|_{A_o + iB_o} \equiv c = c_1 + ic_2 \quad (13)$$

⁴Strictly speaking, our derivation of ω thus far applies only in the limit of global supersymmetry. In subsection 3.2 we will explicitly verify that ω is indeed the superpotential.

In general this constant⁵ is complex as we have emphasised in the above formula. In particular the real part is only well defined modulo 1 (in appropriate units) and is essentially the more familiar real Chern-Simons invariant. Its imaginary part however can in general be any real number. In particular c_2 can - as we will demonstrate - be large. This fact will be important for us in the following.

Summarising the results of this subsection: the contribution of the fields supported at the singularities to the superpotential is given by a complex number, c - the Chern-Simons invariant of a flat complex connection.

3. Minima of the Supergravity Lagrangian.

Having identified the contribution to the superpotential we can add it to the G -flux induced linear superpotential and study the minima of the supergravity potential. The absolute minima of the potential are supersymmetric vacua with a negative cosmological constant. These vacua are characterised by solutions to the following equations:

$$\frac{\partial W}{\partial Z_k} + \frac{\partial K}{\partial Z_k} W = 0 \quad (14)$$

With W given by⁶

$$W = \frac{1}{8\pi^2} (\Sigma_j Z_j G_j + c_1 + i c_2) \quad (15)$$

and the first derivative of the Kahler potential by [30]

$$\frac{\partial K}{\partial Z_k} = \frac{7}{2} i \frac{\int_X \phi_k \wedge * \varphi}{\int_X \varphi \wedge * \varphi} \quad (16)$$

we see that minima exist if,

$$\Sigma t^j G_j + c_1 = 0 \quad (17)$$

and

$$G_k = \frac{7}{2} \frac{\int_X \phi_k \wedge * \varphi}{\int_X \varphi \wedge * \varphi} (\Sigma_j s_j G_j + c_2) \quad (18)$$

3.1. Solution to Vacuum Equations.

⁵Note that if we had chosen a more general non-flat connection, then ω will not be constant and the formulae below become more complicated.

⁶We have redefined c_1 and c_2 for convenience

The equation involving the axions implies that one linear combination of axions has a fixed vev. Note that in the quantum theory this equation gets modified by the addition of half an integer, determined by which “theta vacuum” is picked.

The equations for the moduli fields s_i imply that in any given solution the 4-form field G is proportional to $*\varphi$, with a coefficient α which depends upon the moduli. To see this explicitly, expand $*\varphi$ in terms of the basis of harmonic 4-forms ρ_i which are dual to the ϕ_i :

$$*\varphi = \sum_j u_j \rho_j \quad (19)$$

Note that the u_j depend upon the moduli fields s_i . The u_j are homogeneous of degree $\frac{4}{3}$. Then the equations above may be rewritten as

$$G_k = \frac{7}{2} \frac{u_k}{\sum_j s_j u_j} (\sum_i s_i G_i + c_2) \quad (20)$$

Therefore,

$$G = \alpha * \varphi \quad (21)$$

where

$$\alpha = \frac{7}{2 \sum_j s_j u_j} (\sum_i s_i G_i + c_2) = \frac{1}{2 \text{Vol}(X)} (\sum_i s_i G_i + c_2) \quad (22)$$

Since a solution of the equations implies that $*\varphi$ is proportional to an integral cohomology class, any solution has at most one modulus - the volume. This is because changes in the moduli which preserve the volume of X will always change the direction in $H^4(X, \mathbf{R})$ in which $*\varphi$ points. We will now describe a way to solve these equations and demonstrate that the volume modulus is also fixed.

The idea is simple and we are indebted to Dominic Joyce for proposing it. We assume that our G_2 -manifold X has a G_2 -structure φ' which is proportional to the flux G ,

$$\alpha' * \varphi' = G \quad (23)$$

Then we simply rescale the moduli of this G_2 -metric and try to solve equation (21). Note that if X does not have a G_2 -structure for which (23) is true for some fixed G -flux, then we will typically be able to change the flux and adjust the moduli so that it is true. Hence we can typically find combinations of G_2 -manifolds and fluxes for which (23) is true.

Under a rescaling of the moduli,

$$\varphi' \longrightarrow \lambda^3 \varphi' \quad (24)$$

$$\varphi' \longrightarrow \lambda^4 * \varphi' \quad (25)$$

$$\alpha' \longrightarrow \lambda^{-4} \alpha' \quad (26)$$

Thus, denoting the moduli in the φ' metric as s'_j , in order to solve our equations we have that

$$s_j = \lambda^3 s'_j \quad (27)$$

and require that

$$\alpha = \lambda^{-4} \alpha' \quad (28)$$

This gives the following equation for the scale factor

$$\lambda^3 \Sigma_j s'_j G_j + c_2 = 2\lambda^3 \text{Vol}(X') \alpha' \quad (29)$$

This has a unique solution given by

$$\lambda^3 = \frac{c_2}{2\text{Vol}(X') \alpha' - \Sigma_j s'_j G_j} \quad (30)$$

Hence this minimum of the potential is isolated. In fact it is actually clear from the conditions for unbroken supersymmetry (18) that the volume modulus will be fixed in a given solution. This is because the equations are not invariant under a rescaling of the moduli fields. The left hand side and the first term on the right are invariant, but the last term involving c_2 is not invariant.

In summary, we have shown that if X admits a G_2 -metric which is such that $*\varphi'$ is proportional to G then there exists an isolated minimum of the supergravity potential with negative cosmological constant. Note that X must be such that the topology of Q is rich enough to admit flat connections with non-zero c_2 . We will discuss this issue later. For the moment we wish to outline a dual Type IIA description of the moduli fixing mechanism ⁷ which may serve to clarify the origin of c_2 .

3.2 Type IIA dual description.

In addition to being dual to compactifications of the heterotic string theory, G_2 -compactifications of M theory can also be dual to orientifolds of Type IIA theory on a Calabi-Yau threefold Z . Such string vacua can be described in the following way. Consider Type IIA on a Calabi-Yau threefold Z . This theory has $\mathcal{N} = 2$ supersymmetry. If Z admits a $\mathbf{Z}_2$ symmetry σ under which the Kahler form K and holomorphic 3-form Ω behave as

$$K \rightarrow -K \quad (31)$$

$$\Omega \rightarrow \Omega^* \quad (32)$$

⁷We thank E. Witten for suggesting this.

then one can define an orientifold of the theory by gauging the $\mathbf{Z}_2$ symmetry generated by the product of σ and the world-sheet parity operator P . Gauging this $\mathbf{Z}_2$ breaks the supersymmetry to $\mathcal{N} = 1$.

The fixed points of Z under σ , if non-zero, are 3-dimensional submanifolds Z^σ of Z and will be smooth if Z is smooth. The string theory thus has orientifold 6-planes with topology $Z^\sigma \times M^{3,1}$, with M four-dimensional spacetime. Consequently, the theory also has a tadpole for the RR 7-form potential C_7 , supported on the orientifold planes. The tadpoles may be cancelled by introducing $D6$ -branes wrapping 3-dimensional submanifolds Q_a of Z . If N^a denotes the number of branes wrapping Q_a , the homology classes of the Q_a 's must be such that

$$\Sigma_a N^a [Q_a] = 4[Z^\sigma] \quad (33)$$

This is the tadpole cancellation condition. Some examples of such vacua have been described in [36].

If all the Q_a are special Lagrangian 3-manifolds (ie supersymmetric cycles) then the configuration of branes and orientifolds will be supersymmetric. Since such Type IIA vacua have only the metric, dilaton and RR 1-form potential (dual to C_7) as the only non-zero fields, they arise in the M theory limit solely from the eleven dimensional metric. Supersymmetry then implies that the M theory limit is a compactification on a G_2 -manifold. For instance, the M theory limit of N^a $D6$ -branes on Q_a is an A_{N_a-1} -singularity over Q_a in a G_2 -manifold.

The description of the fields on the $D6$ -branes at low energies is precisely the 7-dimensional super Yang-Mills theory we discussed in the M theory limit. The gauge field A on Q corresponds to the usual gauge field of the $D6$ -brane and the 1-form B on Q has three components which locally represent normal fluctuations of the brane inside the Calabi-Yau Z . The superpotential from the $D6$ -brane sector is again given by the complex Chern-Simons functional, but now we can interpret contributions to its imaginary part as being associated with non-zero values for the normal bundle fields B , since the imaginary part vanishes if B vanishes.

What kind of brane physics is associated with non-zero normal bundle fields? One natural guess is the Myers effect [37]. For example, a non-zero value for

$$tr B \wedge B \wedge B \quad (34)$$

on a Dp -brane induces a coupling to the RR $(p+4)$ -form field strength F of the form

$$tr B^1 B^2 B^3 . F + \text{permutations} \quad (35)$$

where the $(p+4)$ -form F is contracted with the three components of B to give a $(p+1)$ -form which we may integrate over the Dp -brane world-volume. Since the $(p+4)$ -form measures the charge distribution of $D(p+2)$ -branes in Type IIA/B theory, this coupling is associated with Dp -branes which have “expanded into” $D(p+2)$ -branes. In our context $p = 6$ and the non-zero value of (34) occurs when $c_2 \neq 0$. Hence the $D6$ -branes of the orientifold compactification expand into $D8$ -branes.

3.2 The Potential.

Up to now, we have described the conditions for unbroken supersymmetry and their solutions. Also of interest is the supergravity potential, V . V is determined by the standard formula

$$V = e^K (g^{i\bar{j}} (\frac{\partial W}{\partial Z_i} + \frac{\partial K}{\partial Z_i} W) (\frac{\partial \bar{W}}{\partial \bar{Z}_{\bar{j}}} + \frac{\partial K}{\partial \bar{Z}_{\bar{j}}} \bar{W}) - 3|W|^2) \quad (36)$$

This turns out to be given by

$$V = \frac{\pi^2}{2} \left(\frac{\int_X G_X \wedge *G_X}{Vol(X)^2} + \frac{(\int_X \frac{C}{2} \wedge G_X + c_1)^2}{Vol(X)^3} + \frac{(c_2)^2}{Vol(X)^3} + \frac{c_2 \int_X \Phi \wedge G_X}{Vol(X)^3} \right) \quad (37)$$

Note that c_1 and $\int_X \frac{C}{2} \wedge G$ appear on an equal footing in the effective potential. We can see this explicitly from the C -field equations of motion in the presence of the ADE -singularity supported on $Y = Q \times M^{3,1}$. In the presence of such an ADE singularity the M theory Lagrangian has a contribution coupling the gauge fields on Y to the C -field:

$$\Delta L = \delta_Y^4 \wedge \omega_3(A) \wedge G \quad (38)$$

where ω_3 is the Chern-Simons 3-form as a functional of the gauge field and δ_Y^4 is a 4-form delta-function supported on Y . This means that the C -field equation of motion is

$$d * G + \frac{1}{2} G \wedge G + \delta_Y^4 \wedge tr F \wedge F = 0 \quad (39)$$

This implies that the gauge field makes a contribution to the conserved Page charge, P , defined by [38]

$$P = \int_X (*G + \frac{1}{2} C \wedge G) / 4\pi^2 + c_1 \quad (40)$$

Hence, turning on A is equivalent to shifting the Page charge by the Chern-Simons invariant c_1 . We can also use this analysis to reconfirm our calculation

of W using the fact that *BPS* domain walls in the supergravity theory have a tension which is proportional to the change in W as the wall is crossed.

Consider an $M2$ -brane whose world-volume is $q \times \mathbf{R}^{2,1} \subset Q \times \mathbf{R}^{3,1}$. From the effective field theory point of view this domain wall in four dimensions can be described as an instanton on $Q \times \mathbf{R}$ for the gauge field A . Here $\mathbf{R}$ is the direction transverse to the wall. The reason for this is precisely the equation of motion (39) which implies that gauge field configurations of non-zero instanton number have non-zero membrane charge⁸.

To specify the instanton field requires boundary conditions at $\pm\infty$ which consist of two copies of Q . The requisite boundary conditions are a flat connection on each copy of Q . The $M2$ -brane domain wall is then an instanton which interpolates between the two flat connections at the two ends. The tension of such a domain wall is then given by evaluating the Yang-Mills action on $Q \times \mathbf{R}$

$$T = \int_{Q \times \mathbf{R}} \text{tr} F \wedge *F = \int_{Q \times \mathbf{R}} \text{tr} F \wedge F = c_1^+ - c_1^- \equiv \Delta c_1 \quad (41)$$

ie is the difference in the Chern-Simons invariants of the two flat connections. Since the supersymmetry algebra implies that

$$T = \Delta(e^{\frac{K}{2}} W) \quad (42)$$

we find (by comparing with the formulae for V) that W receives a contribution which is directly proportional to c_1 , since $e^K \propto \text{Vol}(X)^{-3}$. Holomorphy ie the fact that the gauge field on Q naturally gets complexified, then implies the c_2 contribution. This completes the derivation of W that we have used throughout this paper.

An open question is what kinds of domain walls in the theory shift the value of c_2 ?? The answer appears to be domain walls which couple only to the gravitational field. This is because the C -field equation of motion does not detect changes in c_2 and therefore the Einstein equations presumably do.

3.4. The One Modulus Case.

We have shown above that the equations formally have an isolated minimum. However, for consistency, we must also show that the minimum exists in a region of field space where the supergravity approximation is valid, since this after all is where we obtained the Kahler potential. This requires showing that the minimum exists at values of the s_j for which the size of X is

⁸In the Type IIA description the instanton is a D2-brane inside the D6-brane.

large compared to the Planck length. Although this is a straightforward exercise in the general case we will restrict our attention to the very simple case of G_2 -manifolds with one modulus, s . In this case we can be very explicit about the dependence of the equations on s and the formulae are a little less abstract.

With one modulus, homogeneity of e^K fixes the Kahler potential to be

$$K = -\ln\left(\frac{1}{14\pi^2}s^7\right) \quad (43)$$

Then the vacuum equations assert that

$$G = \frac{7}{2s}(Gs + c_2) \quad (44)$$

This has the unique solution

$$s = -\frac{7}{5}\frac{c_2}{G} \quad (45)$$

Note that, since s is positive in the supergravity approximation, the signs of c_2 and G must be opposite. Since c_2 is fixed by the topology there is a solution for only one sign of G .

The supergravity approximation is valid when the volume of X is large and this corresponds to s being large. Therefore as long as c_2 is large compared to G , the minimum exists in a region of field space within the approximation. In M theory s has length dimension three, so a c_2 of order 10^3 or bigger and a minimal flux quantum is presumably enough to put the minimum in the regime of validity. Before we discuss in more detail the nature of c_2 and the existence of G_2 -manifolds with non-zero c_2 we can compute using the supergravity potential the value of the potential in the minimum. This is the cosmological constant of the anti-de Sitter space. This turns out to be

$$V_0 = \Lambda_0 = \frac{21\pi^2 5^5}{2 \times 7^5} \left(\frac{G}{c_2}\right)^5 G^2 \approx 20 \frac{G^7}{c_2^5} \quad (46)$$

which if G is minimal is small for large c_2 .

The potential also has another critical point at finite s . This critical point is a de Sitter maximum. The cosmological constant of de Sitter space is of the same order as it is in the minimum. The multi modulus case also presumably has a de Sitter maximum. This follows simply from the fact that, as we have seen, the potential has a minimum below zero at some finite value of the s -fields. But for extremely large values of the fields the terms involving c_2 are negligible compared to the flux induced terms calculated in [30]. These

terms are purely positive and for large s_i the potential approaches zero from above. Hence, presuming the potential does not blow up at finite distance, it must have a maximum above zero. In the multi-modulus case we actually expect the existence of de Sitter minima as well although we have not been able to prove this. This is simply because there are in general many fields. If this is the case, the theory would have non-supersymmetric metastable vacua with positive cosmological constant. Aspects of cosmology of these compactifications are being investigated in [34].

We now turn to the question of the existence of G_2 -manifolds with the properties required for this moduli fixing mechanism to be implemented.

4. On the Existence of G_2 -manifolds with $c_2 \neq 0$.

Recall that our G_2 -manifold X has an ADE -singularity supported along a 3-manifold Q . c_2 is the imaginary part of the Chern-Simons invariant of the flat complex ADE -connection on Q . In particular if B , the imaginary part of the gauge field is gauge equivalent to zero c_2 will vanish. For example, consider the case when the gauge group is $G^{\mathbf{C}} = SL(N, \mathbf{C})$. The flat connection is then a set of matrices in $G^{\mathbf{C}}$ which generate a group which is homomorphic to $\pi_1(Q)$. If, by conjugation by elements in $G^{\mathbf{C}}$, one can unitarise these matrices ie conjugate them into $G^{\mathbf{R}} = SU(n)$, B is gauge equivalent to zero and c_2 vanishes. This is certainly the case when $\pi_1(Q)$ is finite. We should therefore consider the case of infinite fundamental group.

As far as we are aware, the only known examples of flat connections with non-zero c_2 are in the cases in which,

$$Q = \mathbf{H}^3/\Gamma \tag{47}$$

ie when Q is diffeomorphic to a compact 3-manifold which admits a hyperbolic metric⁹. A hyperbolic metric is a metric of constant negative curvature ie is locally isometric to Euclidean anti-de Sitter space.

Now, since

$$\mathbf{H}^3 = SO(1, 3)/SO(3) = PSL(2, \mathbf{C})/PSU(2) \tag{48}$$

Γ is therefore a subgroup of $PSL(2, \mathbf{C})$ and canonically defines a representation of $\pi_1(Q)$ in $PSL(2, \mathbf{C})$ ie a flat $PSL(2, \mathbf{C})$ connection on Q . This representation can always be lifted to a representation of $SL(2, \mathbf{C})$. This is the flat connection that we will consider using, by regarding $SL(2, \mathbf{C})$ as a subgroup of $G^{\mathbf{C}}$. The corresponding gauge field in the standard hyperbolic metric is given by

⁹In [39], a link between G_2 -manifolds and hyperbolic geometry was also made.

$$A + iB = \begin{pmatrix} f & g \\ h & -f \end{pmatrix} \quad (49)$$

where f, g, h are complex one forms given in terms of the dreibein e_i and spin connection ω_{ij} by [40]

$$2f = ie_1 - \omega_{23} \quad (50)$$

$$2g = i(e_2 + \omega_{12}) + e_3 + \omega_{13} \quad (51)$$

$$2h = i(e_2 - \omega_{12}) - e_3 + \omega_{13} \quad (52)$$

Then, one can readily compute c the Chern-Simons invariant in the hyperbolic metric to learn that

$$c_2 \propto Vol(\mathbf{H}^3/\Gamma) \quad (53)$$

So in this case c_2 is the volume of Q computed in its canonically normalised hyperbolic metric. c_2 can in fact be large in practise. To see this, consider a hyperbolic 3-manifold Q' with a volume invariant V' . Then Q' admits a finite cover Q whose volume invariant is NV' . N can be an integer of arbitrarily high order!

The existence of such a Γ implies that Γ contains a normal subgroup of index N . Equivalently, Γ is homomorphic to a finite group of order N .

To show this recall that Γ is finitely generated by determinant one complex matrices (modulo ± 1). The entries of these matrices are complex numbers. Let $\mathbf{F} \subset \mathbf{C}$ denote the sub-ring of the complex numbers generated by the entries of these matrices. $\mathbf{F}$ is also finitely generated. In fact Γ is conjugate to a representation in which the entries of the matrices are all algebraic¹⁰[41]. This implies that Γ is conjugate to $PSL(2, \mathbf{F})$, for some $\mathbf{F}$.

For instance $\mathbf{F}$ might be a subset of $\mathbf{C}$ of the form $\{a + ib | a, b \in \mathbf{Z}\}$. We can then consider the subrings of $\mathbf{F}$ called “prime ideals” which we denote by $\mathbf{P}$. For example, $\mathbf{P}$ might be $\{cp + idp | c, d \in \mathbf{Z}\}$ with p a prime number. Then $\mathbf{F}/\mathbf{P}$ is a finite field and we have maps

$$\Gamma \longrightarrow PSL(2, \mathbf{F}) \longrightarrow PSL(2, \mathbf{F}/\mathbf{P}) \quad (54)$$

This gives a map from Γ to a finite group $PSL(2, \mathbf{F}/\mathbf{P})$ which by choosing the “prime number” in $\mathbf{P}$ is of arbitrarily large order. The kernel of this map is then the fundamental group of a hyperbolic 3-manifold whose volume is of arbitrary order.

¹⁰ie contained in a finite algebraic extension of the field of rational numbers.

Note that if a G_2 -manifold has an ADE -singularity supported along a 3-manifold Q of hyperbolic type then the metric induced on Q will typically not be the constant curvature metric. However, a flat connection with the same Chern-Simons invariant exists on Q regardless of the metric induced upon it. Therefore c_2 is not the actual volume of Q which in general depends upon the s_j . However, because Q is a supersymmetric 3-cycle its volume form is given by φ . Hence the volume of Q is linear in the s_j . Formula (45) (and its generalisation to the multi-modulus case) then shows that the actual volume of Q and its volume in the hyperbolic metric are linearly related.

Another important point about the canonical flat connection on a hyperbolic 3-manifold is that it is rigid ie that it admits no deformations preserving flatness. This was another property that we were assuming in the calculation of the minimum of the effective potential. This property follows from the uniqueness property of hyperbolic 3-manifolds called Mostow Rigidity. The algebraic version of Mostow rigidity asserts that any two isomorphic subgroups of $PSL(2, \mathbb{C})$ which occur as the fundamental groups of finite volume hyperbolic three-manifolds are conjugate. This implies that the flat connection is unique¹¹.

In summary, we have seen that if the G_2 -manifold has an ADE -singularity supported along a 3-manifold Q which admits a hyperbolic metric then there exists an isolated connection with non-zero c_2 . Moreover c_2 can be a large number. Do such G_2 -manifolds exist?

An important result which emerged from Thurston's work on 3-manifolds is that "most" compact 3-manifolds are of hyperbolic type. The Thurston program [41] resulted in a picture of a 3-manifold in which it can be decomposed into a collection of 'prime pieces' which consist of other 3-manifolds. Each prime 3-manifold then admits one of eight types of locally homogeneous metrics. Seven of these eight types are 3-manifolds which are classified. For example they include finite quotients of S^3 and T^3 . The remaining geometry, namely hyperbolic 3-space, makes up all of the remaining prime 3-manifolds and since the list of the other 7-types is so small, hyperbolic 3-manifolds make up "most" prime 3-manifolds.

A more concrete argument pertaining to the existence of hyperbolic submanifolds of G_2 -manifolds is perhaps the following. From various string duality [42, 8] and mathematical arguments [31], one expectation about some G_2 -manifolds is that they admit $K3$ -fibrations over $\mathbf{S}^3$ (or a quotient). Imagine that the generic $K3$ -fiber has not one but several ADE -singularities of the same type. For example three A_2 -singularities. Then locally, each such singularity, as we follow it around on $\mathbf{S}^3$, gives a section of the fibration.

¹¹The proof of Mostow rigidity is somewhat outside the scope of this paper.

However, along a codimension two locus L in $\mathbf{S}^3$ the three singularities of the $K3$ -fibers above L could meet. In this way the $K3$ -fibration would not have a section but an order 3 multi-section. This produces in the total space of X a single A_2 -singularity supported on a 3-manifold Σ . Σ is in fact a 3-fold branched cover of $\mathbf{S}^3$. An important fact about 3-manifolds is that any 3-manifold can be regarded as a 3-fold branched cover of $\mathbf{S}^3$ [43, 44]. So, in this way one can imagine that X is such that the branching is hyperbolic. Unfortunately, developing this argument further requires understanding better the branching locus¹² L .

We should also point out an analagous situation in which plenty of examples may be found. This consists of a compact Calabi-Yau threefold Z with an ADE -singularity localised along a hyperbolic 2-manifold ie a Riemann surface of genus > 1 .

Having given some arguments which might suggest that G_2 -manifolds exist with non-zero c_2 we now return to the physics - assuming the existence of a solution.

5. Holography and AdS/CFT.

The alert reader may have noticed something which at first sight appears peculiar from our discussion of the minima of the supergravity potential. This is the fact that the potential for the s_i had an isolated minimum but only one linear combination of axions had a fixed vev in the vacuum. It thus appears that we have an $\mathcal{N} = 1$ vacuum configuration in which $b_3(X) - 1$ real scalars (the axions) are massless. This seems peculiar at first because the usual representations of the $\mathcal{N} = 1$ supersymmetry algebra contain two bosonic degrees of freedom not one. As we shall explain this appears to be related to the holographic dual description of these vacua.

The resolution of this peculiarity is that the vacuum we are talking about is a supersymmetric anti-de Sitter vacuum. In anti-de Sitter space, in addition to supermultiplets with two bosonic degrees of freedom, there does indeed exist a supermultiplet with one bosonic degree of freedom. This is the so called singleton representation. Therefore, our axions must be singletons.

An important point about singleton representations is that they naturally reside on the boundary of anti-de Sitter space¹³. Thus, the excitations of the axion fields in our vacua represent boundary degrees of freedom.

¹² X could have an additional singularity along L at which additional light degrees of freedom might be found.

¹³See the introduction of [45] for a review with detailed references on the representation theory of the anti-de Sitter group.

A very natural question to ask is what is the holographic dual description of these adS vacua? Because of the symmetries, this dual description is almost certainly a three dimensional conformal field theory which must - in some natural sense - reside on the boundary of anti-de Sitter space. In the better understood adS vacua of M theory, this CFT is the world-volume theory of some collection of $M2$ -branes on the boundary of the spacetime. In that context it was realised that the singleton representation in the gravity theory represented the center of mass degree of freedom of the branes.

Since we have identified $b_3(X) - 1$ axionic singleton fields in the adS vacua under discussion here, it is natural to propose that the holographic dual description corresponds to the world-volume field theory of $b_3(X) - 1$ collections of branes at the boundary of the spacetime (although it is not clear which kind of branes these actually are). One could try and verify this proposal by finding the corresponding brane solution of M theory on X .

6. Phenomenology.

Presuming that compact G_2 -manifolds with non-zero c_2 exist, we have shown that vacua of M theory exist with no moduli. This means that such G_2 -manifolds would provide solutions of the eleven dimensional theory which describe four dimensional universes with no undetermined parameters except the fundamental Planck scale. In principle therefore all masses and couplings are calculable by the theory in the isolated vacuum state.

The main phenomenological difficulties with the vacua we have described here are a) they are supersymmetric and b) they have a negative cosmological constant. However, as we have seen, the moduli fixing mechanism that we have described here can in principle be implemented in a G_2 -manifold whose associated low energy physics contains no light charged matter, since we did not have to consider the matter fields supported at “chiral singularities” of X . We should therefore consider the utility of the mechanism in compactifications which include quarks and leptons.

The basic idea that we consider is that the moduli fixing mechanism considered here is the dominant contribution to the moduli potential in a more realistic scenario. In order to investigate this possibility, it seems necessary to assume that in addition to the singularities which produce a supersymmetric standard model sector, X also has singularities which produce a supersymmetry breaking sector. Once supersymmetry is broken (at a scale m_{susy}) the zero point energies of the charged fields will generate a contribution Λ_1 to the cosmological constant of order m_{susy}^4 and we will simply consider the consequences of requiring that Λ_1 and the contribution Λ_0 of section three are of the same order. This is necessary if one is to obtain a positive cosmolog-

ical constant. Under the *strong* assumption that a suitable supersymmetry breaking mechanism exists for which this is true, there are a couple of quite striking consequences.

First, the heirarchy problem reduces to explaining why a topological invariant c_2 takes the value that it does. Second, the fundamental Planck scale is roughly 10 TeV, if we take $m_{susy} = 1$ Tev. Thus, if these sorts of models actually existed, they would provide a concrete realisation of the ideas proposed in [46].

Note that the cosmological constant problem in this context then reduces to the problem of explaining why Λ_1 cancels Λ_0 to at least 10^{56} orders of magnitude. We now turn to an explanation of these statements.

Once supersymmetry is broken, the mass splittings between fermions and bosons will generate a contribution to the cosmological constant of the form m_{susy}^4 , where m_{susy} is the scale of supersymmetry breaking. If this contribution to Λ is going to be comparable to the cosmological constant calculated in section three, we require that

$$20 \frac{G^7}{c_2^5} m_p^4 \approx m_{susy}^4 \quad (55)$$

where m_p is the four dimensional Planck mass. Thus, the heirarchy problem essentially reduces to explaining why

$$\frac{c_2^5}{G^7} \approx 20 \left(\frac{m_p}{m_{susy}} \right)^4 \quad (56)$$

The minimal value for G is 2π , so the smallest c_2 can be is around $10^{13} - 10^{14}$, if we assume that $1TeV \leq m_{susy} \leq 10TeV$. So in this scenario, the heirarchy problem reduces to explaining why a certain topological invariant is 10^{13} (or bigger if the fluxes are larger).

On the other hand, m_p is determined in terms of the eleven dimensional Planck mass, M_p through the formula

$$m_p^2 \approx \left(\frac{c_2}{G} \right)^{7/3} M_p^2 \quad (57)$$

Hence, assuming that m_{susy} is around the TeV scale and we know m_p , we can determine the fundamental Planck scale. For the minimal value of c_2 , we find that the fundamental scale is one order larger than m_{susy}

$$M_p \approx (10 - 100)TeV \approx 10m_{susy} \quad (58)$$

For a much larger G -flux, say of order 10^{10} ,

$$c_2 \approx 10^{27} \quad (59)$$

and

$$M_p \approx 1\text{GeV} \quad (60)$$

so vacua in which c_2 and G are vastly different, have fundamental scales which differ by a much milder amount. However, a fundamental scale which is lower than the scales already probed by particle physics experiments is surely problematic and hence models with flux-quanta which are too large are ruled out.

We should also compute the masses of Kaluza-Klein particles. These are given by

$$m_{kk} \approx \left(\frac{G}{c_2}\right)^{1/3} M_p \quad (61)$$

For minimal G -flux, this is roughly 10 GeV. For larger fluxes these masses are smaller.

An important comment is in order. If X is isotropic in the minimum all its moduli are of the same order of magnitude in the minimum of the potential. Naively this would imply that these vacua are ruled out by accelerator experiments, because Kaluza-Klein excitations of the gauge fields along the Q_a which supports the standard model gauge group would produce charged particles of mass 10 GeV or less. Renormalisable couplings between these particles and those of the standard model would then imply that such isotropic vacua are inconsistent else the Kaluza-Klein modes would have been observed in particle physics experiments above 10 GeV.

However, recall that the standard model particles are supported at singular points on X and therefore transform only under four-dimensional gauge transformations, whereas the gauge fields transform under *seven* dimensional gauge transformations. The charged Kaluza-Klein modes therefore transform under seven dimensional gauge transformations. Because of this, *renormalisable* interactions between standard model particles and the Kaluza-Klein modes are difficult to imagine and hence the interactions of these modes will presumably be much harder to detect.

Even if the charged Kaluza-Klein modes do have conventional couplings to quarks and leptons, many models could exist in which their masses are 100 GeV or much larger. To see this, we consider the generalisation of (45) to the general case,

$$\Sigma_i s^i G_i = -\frac{7c_2}{5} \quad (62)$$

The Q 's which support the gauge fields have volumes which are linear in the s^i because they are supersymmetric cycles. Hence if s^1 is the volume of one of the Q 's the above formula asserts that as G_1 is increased, s^1 decreases. So a larger flux in one particular direction in the cohomology of X

corresponds to shrinking a 3-manifold in X . Therefore, the Q which supports the standard model gauge group can easily be much smaller than the largest scale of X and if the mass scale of Q is 100 GeV or bigger the model is consistent with accelerator experiments. These smaller 3-cycles make a negligible contribution to the volume of X . This implies that the cosmological constant of section three is still given by (46), but with the G which appears there the smallest component of G -flux in X .

It would certainly be of interest to develop some of the ideas presented here further. In particular, a construction of G_2 -manifolds with some of the properties described here would be an important step.

Appendix: Supersymmetry.

A relatively efficient method to derive the equations for unbroken supersymmetry on the fields supported at the ADE -singularity along Q is actually in a dual context. Namely when Q is a supersymmetric 3-cycle in a Calabi-Yau threefold Z in Type IIA and some number of $D6$ -branes are wrapping Q . Then, as shown in [3], the supersymmetry conditions for the world-volume fields are identical to those in M theory. This is because the strong coupling limit of this Type IIA configuration is precisely an ADE -singularity along Q .

In flat space the world-volume theory of N $D6$ -branes is $U(N)$ super Yang-Mills in seven dimensions. We can derive the conditions for unbroken supersymmetry by dimensional reduction of the corresponding conditions in ten dimensional super Yang-Mills theory on a $D9$ -brane. In curved space, the analogous procedure is to dimensionally reduce the conditions for unbroken supersymmetry for gauge fields along Z to Q . Namely, we should consider the dimensional reduction of the Hermitian Yang-Mills equations on Z onto Q .

In components in a locally flat frame, the Hermitian Yang-Mills equations are linear equations on the field strength F_{AB} of a gauge field a_B on Z :

$$F_{12} + F_{34} + F_{56} = 0 \tag{63}$$

$$F_{13} - F_{24} = 0 \tag{64}$$

$$F_{14} + F_{23} = 0 \tag{65}$$

$$F_{15} - F_{26} = 0 \tag{66}$$

$$F_{16} + F_{25} = 0 \tag{67}$$

$$F_{35} - F_{46} = 0 \tag{68}$$

$$F_{36} + F_{45} = 0 \tag{69}$$

$$\tag{70}$$

The first equation depends upon the metric on Z via the Kahler form which we have chosen to be

$$\omega = dx_1 \wedge dx_2 + dx_3 \wedge dx_4 + dx_5 \wedge dx_6 \quad (71)$$

The remaining six equations are the conditions that the gauge field is holomorphic. These do not depend upon the Kahler metric. These equations correspond to F -terms in four dimensions and are the critical points of the holomorphic Chern-Simons functional.

Q is a supersymmetric 3-cycle in Z and as such

$$\omega|_Q = 0 \quad (72)$$

So in the local frame e_i we will choose the frames along Q to be spanned by (e_1, e_3, e_5) . The $\mathbf{R}^3$ -subspace of $\mathbf{C}^3$ thus defined is indeed special Lagrangian with the complex structure defined by ω . Correspondingly we will define the gauge field components tangential and normal to Q by

$$(a_1, a_3, a_5) \equiv (A_1, A_2, A_3) \quad (73)$$

$$(a_2, a_4, a_6) \equiv (B_1, B_2, B_3) \quad (74)$$

Dimensionally reducing the Hermitian Yang-Mills equations onto Q then corresponds to setting to zero the normal derivatives:

$$\partial_2 = \partial_4 = \partial_6 = 0 \quad (75)$$

If we regard both A_i and B_i as components of a 1-form on Q (with values in the Lie algebra of the gauge group) then the above seven equations become:

$$d * B + A \wedge * B + * B \wedge A = 0 \quad (76)$$

$$dA + A \wedge A = B \wedge B \quad (77)$$

$$dB + A \wedge B - B \wedge A = 0 \quad (78)$$

The first of these depends upon the metric on Q and corresponds in four dimensions to a D -term. The remaining six equations do not depend upon the metric and are F -terms. They are the critical points of a complex Chern-Simons action, as described in the text.

Acknowledgements.

We would like to thank M. Atiyah, T. Banks, M. Douglas, N. Dunfield, F. Denef, D. Friedan, D. Joyce, S. Kachru, N. Lambert, D. Long, J. Maldacena, G. Moore, E. Silverstein and E. Witten for very useful discussions.

References

- [1] M. Dine and N. Seiberg, Is The Superstring Semiclassical?, *Unified String Theories: Proceedings* eds. M.B. Green and D. Gross, World Scientific, 1986.
- [2] R. Brustein and P. Steinhardt, Challenges For Superstring Cosmology, *Phys. Lett. B* **302**, 196 (1993) [arXiv:hep-th/9212049].
- [3] B.S. Acharya, M theory, Joyce Orbifolds and Super Yang-Mills, [arXiv:hep-th/9812205].
- [4] B.S. Acharya, On Realising N=1 super Yang-Mills in M theory. [arXiv:hep-th/0011089]
- [5] M. Atiyah, J. Maldacena and C. Vafa, An M theory flop as a Large N duality, [arXiv:hep-th/0011256].
- [6] M. Atiyah and E. Witten, M theory Dynamics on a Manifold of G_2 -holonomy, [arXiv:hep-th/0107177].
- [7] E. Witten, Anomaly Cancellation on G_2 -manifolds, [arXiv:hep-th/0108165].
- [8] B.S. Acharya and E. Witten, Chiral Fermions from Manifolds of G_2 -holonomy, [arXiv:hep-th/0109152].
- [9] P. Berglund and A. Brandhuber, Matter from G_2 -manifolds, [arXiv:hep-th/0205184].
- [10] E. Witten, Deconstruction, G_2 -holonomy and Doublet-Triplet Splitting, [arXiv:hep-ph/0201018].
- [11] T. Friedmann and E. Witten, Unification Scale, Proton Decay and Manifolds of G_2 -holonomy, [arXiv:hep-th/0211269].
- [12] K. Becker and M. Becker, M-Theory on Eight-Manifolds, *Nucl. Phys. B* **477**, 155 (1996) [arXiv:hep-th/9605053].
- [13] S. Gukov, C. Vafa and E. Witten, CFT's from Calabi-Yau four-folds, *Nucl. Phys. B* **584**, 69 (2000) [Erratum-ibid. *B* **608**, 477 (2001)] [arXiv:hep-th/9906070].
- [14] K. Dasgupta, G. Rajesh and S. Sethi, M theory, orientifolds and G-flux, *JHEP* **9908**, 023 (1999) [arXiv:hep-th/9908088].

- [15] S. Gukov, Solitons, superpotentials and calibrations, Nucl. Phys. B **574**, 169 (2000) [arXiv:hep-th/9911011].
- [16] G. Curio, A. Klemm, D. Lust and S. Theisen, On the vacuum structure of type II string compactifications on Calabi-Yau spaces with H-fluxes, Nucl. Phys. B **609**, 3 (2001) [arXiv:hep-th/0012213].
- [17] M. Haack and J. Louis, M-theory compactified on Calabi-Yau fourfolds with background flux, Phys. Lett. B **507**, 296 (2001) [arXiv:hep-th/0103068].
- [18] G. Curio, A. Klemm, B. Kors and D. Lust, Fluxes in heterotic and type II string compactifications, Nucl. Phys. B **620**, 237 (2002) [arXiv:hep-th/0106155].
- [19] G. Curio and A. Krause, G-fluxes and non-perturbative stabilisation of heterotic M-theory, Nucl. Phys. B **643**, 131 (2002) [arXiv:hep-th/0108220].
- [20] G. Curio, B. Kors and D. Lust, Fluxes and branes in type II vacua and M-theory geometry with G(2) and Spin(7) holonomy, Nucl. Phys. B **636**, 197 (2002) [arXiv:hep-th/0111165].
- [21] S. Kachru, M. B. Schulz and S. Trivedi, Moduli stabilization from fluxes in a simple IIB orientifold, [arXiv:hep-th/0201028].
- [22] B. Acharya, X. de la Ossa and S. Gukov, G-flux, supersymmetry and Spin(7) manifolds, JHEP **0209**, 047 (2002) [arXiv:hep-th/0201227].
- [23] K. Becker, M. Becker, M. Haack and J. Louis, Supersymmetry breaking and α' corrections to flux induced potentials, JHEP **0206**, 060 (2002) [arXiv:hep-th/0204254].
- [24] R. Argurio, V. L. Campos, G. Ferretti and R. Heise, Freezing of moduli with fluxes in three dimensions, Nucl. Phys. B **640**, 351 (2002) [arXiv:hep-th/0205295].
- [25] S. Hellerman, J. McGreevy and B. Williams, Geometric constructions of nongeometric string theories, arXiv:hep-th/0208174.
- [26] S. Gukov and M. Haack, IIA string theory on Calabi-Yau fourfolds with background fluxes, Nucl. Phys. B **639**, 95 (2002) [arXiv:hep-th/0203267].
- [27] S. Ferrara and M. Porrati, $N = 1$ no-scale supergravity from IIB orientifolds, Phys. Lett. B **545**, 411 (2002) [arXiv:hep-th/0207135].

- [28] A. Dabholkar and C. Hull, Duality twists, orbifolds, and fluxes, arXiv:hep-th/0210209.
- [29] B. S. Acharya and B. Spence, Flux, supersymmetry and M theory on 7-manifolds, arXiv:hep-th/0007213.
- [30] C. Beasley and E. Witten, A note on fluxes and superpotentials in M-theory compactifications on manifolds of $G(2)$ holonomy, JHEP **0207**, 046 (2002) [arXiv:hep-th/0203061].
- [31] D.D. Joyce, **Compact Manifolds of Special Holonomy**, Oxford University Press 2001.
- [32] J. Harvey and G. Moore, Superpotentials and Membrane Instantons, [arXiv:hep-th/9907026].
- [33] J. Gutowski and G. Papadopoulos, Brane Solitons for G_2 -structures in Eleven dimensional supergravity re-visited, [arXiv:hep-th/0208051].
- [34] B.S. Acharya, N. Lambert and D. Tong, work in progress.
- [35] E. Witten, New Issues in Manifolds of $SU(3)$ -holonomy. Nucl. Phys. B **268**, 79 (1986).
- [36] M. Cvetič, G. Shiu and A. M. Uranga, Three-family supersymmetric standard like models from intersecting brane worlds, Phys. Rev. Lett. **87**, 201801 (2001) [arXiv:hep-th/0107143]. M. Cvetič, G. Shiu and A. M. Uranga, Chiral four-dimensional $N = 1$ supersymmetric type IIA orientifolds from intersecting D6-branes, Nucl. Phys. B **615**, 3 (2001) [arXiv:hep-th/0107166]. M. Cvetič, P. Langacker and G. Shiu, Phenomenology of a three-family standard-like string model, Phys. Rev. D **66**, 066004 (2002) [arXiv:hep-ph/0205252]. M. Cvetič, P. Langacker and G. Shiu, Phenomenology of a three-family standard-like string model, Phys. Rev. D **66**, 066004 (2002) [arXiv:hep-ph/0205252]. R. Blumenhagen, V. Braun, B. Kors and D. Lust, Orientifolds of K3 and Calabi-Yau manifolds with intersecting D-branes, JHEP **0207**, 026 (2002) [arXiv:hep-th/0206038]. M. Cvetič, P. Langacker and G. Shiu, A three-family standard-like orientifold model: Yukawa couplings and hierarchy, Nucl. Phys. B **642**, 139 (2002) [arXiv:hep-th/0206115]. C. Kokorelis, Deformed intersecting D6-brane GUTs. I, JHEP **0211**, 027 (2002) [arXiv:hep-th/0209202]. R. Blumenhagen, L. Gorlich and T. Ott, Supersymmetric intersecting branes on the type IIA $T^{**6}/Z(4)$ orientifold, arXiv:hep-th/0211059. M. Cvetič, I. Papadimitrou and G. Shiu, Supersymmetric

Three Family SU(5) Grand Unified Models from Type IIA Orientifolds with Intersecting D6-branes, [arXiv:hep-th/0212177].

- [37] R. Myers, Dielectric-branes, JHEP **9912**, 022 (1999) [arXiv:hep-th/9910053].
- [38] D. N. Page, Classical Stability Of Round And Squashed Seven Spheres In Eleven-Dimensional Supergravity, Phys. Rev. D **28**, 2976 (1983).
- [39] G. Curio, Superpotentials for M-theory on a G(2) holonomy manifold and triality symmetry, [arXiv:hep-th/0212211].
- [40] D. Calegari, Chern-Simons Invariants of Hyperbolic Manifolds, Unpublished notes.
- [41] W. P. Thurston, The Geometry and Topology of Three Manifolds, <http://msri.org/publications/books/gt3m>
- [42] B.S. Acharya, N=1 Heterotic/M theory duality and Joyce Manifolds, Nucl. Phys. B **475**, 579 (1996) [arXiv:hep-th/9603033].
- [43] J.M. Montesinos, Three-manifolds as 3-fold branched covers of S^3 , Quart. J. Math. Oxford Ser (2) **27** (1976) 85.
- [44] H. M. Hilden, Every Closed Orientable 3-manifold is a 3-fold branched covering space of S^3 , Bull. Amer. Math. Soc. **80** (1974) 1243.
- [45] M. Gunaydin and D. Minic, Singletons, Doubletons and M theory, Nucl. Phys. **B523** (1998) 145; [arXiv:hep-th/9802047]
- [46] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, The Hierarchy Problem and New Dimensions at a Millimeter, Phys. Lett. **B429** (1998) 263 [arXiv:hep-ph/9803315]; I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, New Dimensions at a Millimeter to a Fermi and Superstrings at a TeV Phys. Lett. **B436** (1998) 257.